



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 21, 2024 – 09:52 AM EDT

PDB ID : 2MLY
BMRB ID : 19836
Title : NMR structure of E. coli Trigger Factor in complex with unfolded PhoA1-150
Authors : Saio, T.; Guan, X.; Rossi, P.; Economou, A.; Kalodimos, C.G.
Deposited on : 2014-03-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 39%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	443	
2	B	151	

2 Ensemble composition and analysis i

This entry contains 10 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:373, A:392-A:429 (411)	1.43	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 4, 5, 6, 7, 8, 9, 10
2	1, 3

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9012 atoms, of which 4519 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Trigger factor.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	432	6789	2119	3403	582	674	11	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP U6N325
A	-9	ASN	-	expression tag	UNP U6N325
A	-8	HIS	-	expression tag	UNP U6N325
A	-7	LYS	-	expression tag	UNP U6N325
A	-6	VAL	-	expression tag	UNP U6N325
A	-5	HIS	-	expression tag	UNP U6N325
A	-4	HIS	-	expression tag	UNP U6N325
A	-3	HIS	-	expression tag	UNP U6N325
A	-2	HIS	-	expression tag	UNP U6N325
A	-1	HIS	-	expression tag	UNP U6N325
A	0	HIS	-	expression tag	UNP U6N325

- Molecule 2 is a protein called Alkaline phosphatase.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	150	2223	694	1116	192	218	3	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP U6N3P1



- Molecule 2: Alkaline phosphatase

Chain B:  99%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 1000 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CYANA	structure solution	3.0
X-PLOR NIH	structure solution	
TALOSN	geometry optimization	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	2
Total number of shifts	2632
Number of shifts mapped to atoms	2632
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	39%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.85±0.01	0±0/3251 (0.0± 0.0%)	0.63±0.01	0±0/4372 (0.0± 0.0%)
All	All	0.85	0/32510 (0.0%)	0.63	2/43720 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	285	ARG	NE-CZ-NH1	5.79	123.20	120.30	5	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	3214	3245	3245	18±5
2	B	0	0	0	0±0
All	All	32140	32450	32450	175

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 98 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:425:ASN:HA	1:A:428:MET:HB2	0.65	1.66	5	3
1:A:132:VAL:HG11	1:A:274:MET:SD	0.60	2.36	4	2
1:A:88:GLY:HA2	1:A:305:ALA:HB2	0.59	1.73	1	1
1:A:137:VAL:HA	1:A:274:MET:HE1	0.58	1.76	4	1
1:A:154:LYS:HG3	1:A:240:VAL:HG13	0.58	1.74	7	9

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	410/443 (93%)	372±5 (91±1%)	36±5 (9±1%)	2±1 (0±0%)	38 78
2	B	0	-	-	-	-
All	All	4100/5940 (69%)	3724 (91%)	361 (9%)	15 (0%)	38 78

5 of 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	326	GLU	2
1	A	246	PRO	2
1	A	149	ALA	2
1	A	95	GLY	2
1	A	12	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	341/370 (92%)	332±2 (97±1%)	9±2 (3±1%)	49 91
2	B	0	-	-	-

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	3410/4840 (70%)	3322 (97%)	88 (3%)	49 91

5 of 29 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	361	LYS	10
1	A	336	LEU	9
1	A	194	MET	8
1	A	236	ASN	7
1	A	294	LEU	5

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 39% for the well-defined parts and 34% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2303
Number of shifts mapped to atoms	2303
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	37

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	91	-0.09 ± 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	106	0.08 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}'$	1	—	None (insufficient data)
^{15}N	392	0.45 ± 0.31	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 39%, i.e. 2224 atoms were assigned a chemical shift out of a possible 5638. 0 out of 72 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	941/2065 (46%)	476/843 (56%)	92/822 (11%)	373/400 (93%)
Sidechain	1172/3331 (35%)	842/2148 (39%)	330/1043 (32%)	0/140 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	111/242 (46%)	44/118 (37%)	66/119 (55%)	1/5 (20%)
Overall	2224/5638 (39%)	1362/3109 (44%)	488/1984 (25%)	374/545 (69%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	337	PHE	CE2	40.88	124.80 – 136.72	-75.4
1	A	322	PHE	CE2	41.09	124.80 – 136.72	-75.2
1	A	252	PHE	CE2	41.30	124.80 – 136.72	-75.0
1	A	256	PHE	CE2	41.42	124.80 – 136.72	-75.0
1	A	387	PHE	CE2	41.43	124.80 – 136.72	-74.9
1	A	424	PHE	CE2	41.95	124.80 – 136.72	-74.5
1	A	256	PHE	CD2	41.95	125.53 – 137.61	-74.2
1	A	387	PHE	CD2	41.95	125.53 – 137.61	-74.2
1	A	424	PHE	CD2	41.95	125.53 – 137.61	-74.2
1	A	252	PHE	CD2	42.05	125.53 – 137.61	-74.1
1	A	337	PHE	CD2	42.22	125.53 – 137.61	-74.0
1	A	322	PHE	CD2	42.79	125.53 – 137.61	-73.5
1	A	378	TYR	CE2	28.15	111.68 – 124.17	-71.9
1	A	388	TYR	CE2	28.19	111.68 – 124.17	-71.8
1	A	256	PHE	CD1	41.95	125.33 – 137.83	-71.7
1	A	387	PHE	CD1	41.95	125.33 – 137.83	-71.7
1	A	424	PHE	CD1	41.95	125.33 – 137.83	-71.7
1	A	252	PHE	CD1	42.05	125.33 – 137.83	-71.6
1	A	337	PHE	CD1	42.22	125.33 – 137.83	-71.5
1	A	322	PHE	CD1	42.79	125.33 – 137.83	-71.0
1	A	337	PHE	CE1	40.88	124.17 – 137.29	-68.5
1	A	322	PHE	CE1	41.09	124.17 – 137.29	-68.3
1	A	252	PHE	CE1	41.30	124.17 – 137.29	-68.2
1	A	256	PHE	CE1	41.42	124.17 – 137.29	-68.1
1	A	387	PHE	CE1	41.43	124.17 – 137.29	-68.1
1	A	424	PHE	CE1	41.95	124.17 – 137.29	-67.7
1	A	378	TYR	CE1	28.15	111.24 – 124.66	-66.9
1	A	388	TYR	CE1	28.19	111.24 – 124.66	-66.9
1	A	388	TYR	CD1	43.26	125.84 – 139.60	-65.0
1	A	378	TYR	CD1	43.67	125.84 – 139.60	-64.7
1	A	337	PHE	CZ	39.15	121.82 – 136.66	-60.7
1	A	322	PHE	CZ	39.16	121.82 – 136.66	-60.7

Continued on next page...


Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	424	PHE	CZ	39.81	121.82 – 136.66	-60.3
1	A	387	PHE	CZ	39.84	121.82 – 136.66	-60.2
1	A	388	TYR	CD2	43.26	125.28 – 140.14	-60.2
1	A	378	TYR	CD2	43.67	125.28 – 140.14	-59.9
1	A	163	ARG	HB2	0.21	0.52 – 3.08	-6.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1_dup*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	329
------------------------	-----

Number of shifts mapped to atoms	329
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	16

7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	4	—	None (insufficient data)
$^{13}\text{C}_\beta$	10	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	84	-0.79 ± 0.19	Should be applied

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 5638. 0 out of 72 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	0/2065 (0%)	0/843 (0%)	0/822 (0%)	0/400 (0%)
Sidechain	0/3331 (0%)	0/2148 (0%)	0/1043 (0%)	0/140 (0%)
Aromatic	0/242 (0%)	0/118 (0%)	0/119 (0%)	0/5 (0%)
Overall	0/5638 (0%)	0/3109 (0%)	0/1984 (0%)	0/545 (0%)

7.2.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
2	B	100	LEU	HD11	25.30	-0.61 – 2.12	89.9
2	B	100	LEU	HD12	25.30	-0.61 – 2.12	89.9
2	B	100	LEU	HD13	25.30	-0.61 – 2.12	89.9
2	B	102	LEU	HD11	25.30	-0.61 – 2.12	89.9
2	B	102	LEU	HD12	25.30	-0.61 – 2.12	89.9
2	B	102	LEU	HD13	25.30	-0.61 – 2.12	89.9

Continued on next page...

Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
2	B	100	LEU	HD21	23.55	-0.65 – 2.13	82.1
2	B	100	LEU	HD22	23.55	-0.65 – 2.13	82.1
2	B	100	LEU	HD23	23.55	-0.65 – 2.13	82.1
2	B	102	LEU	HD21	23.55	-0.65 – 2.13	82.1
2	B	102	LEU	HD22	23.55	-0.65 – 2.13	82.1
2	B	102	LEU	HD23	23.55	-0.65 – 2.13	82.1
2	B	100	LEU	CD1	0.82	16.71 – 32.55	-15.0
2	B	102	LEU	CD1	0.83	16.71 – 32.55	-15.0
2	B	100	LEU	CD2	0.78	15.73 – 32.47	-13.9
2	B	102	LEU	CD2	0.78	15.73 – 32.47	-13.9

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

