



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 4, 2023 – 05:15 PM EDT

PDB ID : 6POY
Title : Structure of human endothelial nitric oxide synthase heme domain in complex with 7-(3-(Aminomethyl)-4-propoxyphenyl)-4-methylquinolin-2-amine
Authors : Chreifi, G.; Li, H.; Poulos, T.L.
Deposited on : 2019-07-05
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 5% 75% 16% 8%
1	B	440	 % 80% 11% 9%
1	C	440	 78% 13% 9%
1	D	440	 77% 14% 8%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 13740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3237	2062	570	589	16	0	2	0
1	B	402	3221	2051	566	587	17	0	3	0
1	C	401	3209	2044	563	586	16	0	2	0
1	D	404	3241	2063	572	589	17	0	3	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	298	GLU	ASP	variant	UNP P29474
B	298	GLU	ASP	variant	UNP P29474
C	298	GLU	ASP	variant	UNP P29474
D	298	GLU	ASP	variant	UNP P29474

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0
2	C	1	43	34	1	4	4	0	0
2	D	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		
3	C	1	Total	C	N	O	0	0
			17	9	5	3		
3	D	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 7-[3-(aminomethyl)-4-propoxyphenyl]-4-methylquinolin-2-amine (three-letter code: OSG) (formula: C₂₀H₂₃N₃O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			24	20	3	1		
4	B	1	Total	C	N	O	0	0
			24	20	3	1		
4	C	1	Total	C	N	O	0	0
			24	20	3	1		
4	D	1	Total	C	N	O	0	0
			24	20	3	1		

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		
6	C	1	Total	Zn	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	C	1	Total C O 6 3 3	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Cl 1 1	0	0
8	B	1	Total Cl 1 1	0	0
8	C	1	Total Cl 1 1	0	0
8	D	1	Total Cl 1 1	0	0

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Gd 1 1	0	0
9	B	1	Total Gd 1 1	0	0
9	C	1	Total Gd 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	1	Total 1	O Gd 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	63	Total 63	O 63	0	0
10	B	105	Total 105	O 105	0	0
10	C	82	Total 82	O 82	0	0
10	D	98	Total 98	O 98	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitric oxide synthase, endothelial




- Molecule 1: Nitric oxide synthase, endothelial




- Molecule 1: Nitric oxide synthase, endothelial





- Molecule 1: Nitric oxide synthase, endothelial

Chain D: 77% 14% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.69Å 153.41Å 109.19Å 90.00° 90.60° 90.00°	Depositor
Resolution (Å)	52.60 – 2.30 88.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.5 (52.60-2.30) 97.3 (88.95-2.30)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, R_{free}	0.198 , 0.261 0.188 , 0.253	Depositor DCC
R_{free} test set	4261 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtrriage
Anisotropy	0.932	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.088 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13740	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, HEM, GD, CL, GOL, H4B, OSG, BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3335	0.55	1/4543 (0.0%)
1	B	0.41	0/3319	0.54	0/4523
1	C	0.40	0/3307	0.53	0/4507
1	D	0.44	0/3339	0.58	0/4548
All	All	0.41	0/13300	0.55	1/18121 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	124	LEU	CA-CB-CG	5.02	126.84	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	238	ARG	Peptide
1	D	89	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3237	0	3146	46	0
1	B	3221	0	3126	33	0
1	C	3209	0	3109	27	0
1	D	3241	0	3152	41	0
2	A	43	0	30	4	0
2	B	43	0	30	4	0
2	C	43	0	30	2	0
2	D	43	0	30	4	0
3	A	17	0	15	1	0
3	B	17	0	15	0	0
3	C	17	0	15	1	0
3	D	17	0	15	1	0
4	A	24	0	0	2	0
4	B	24	0	0	2	0
4	C	24	0	0	1	0
4	D	24	0	0	2	0
5	A	28	0	38	5	0
5	B	42	0	54	10	0
5	C	28	0	37	6	0
5	D	28	0	36	5	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	6	0	8	1	0
7	C	6	0	8	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
10	A	63	0	0	4	0
10	B	105	0	0	5	0
10	C	82	0	0	3	0
10	D	98	0	0	2	0
All	All	13740	0	12894	173	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 173 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:183:ARG:HB2	2:C:501:HEM:HBD2	1.60	0.81
1:D:321:GLU:OE1	5:D:504:BTB:O4	2.08	0.72
1:A:139:LYS:HE3	1:A:140:ARG:HG2	1.71	0.72
1:A:183:ARG:HB2	2:A:501:HEM:O1D	1.90	0.71
1:B:279:TRP:HB2	1:B:302:LEU:HD21	1.73	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/440 (91%)	373 (93%)	24 (6%)	5 (1%)	13	14
1	B	401/440 (91%)	391 (98%)	10 (2%)	0	100	100
1	C	399/440 (91%)	385 (96%)	13 (3%)	1 (0%)	41	50
1	D	403/440 (92%)	389 (96%)	13 (3%)	1 (0%)	47	58
All	All	1605/1760 (91%)	1538 (96%)	60 (4%)	7 (0%)	34	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	283	ASN
1	A	90	GLN
1	C	89	GLN
1	A	145	ALA
1	A	281	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/373 (92%)	333 (96%)	12 (4%)	36	50
1	B	344/373 (92%)	332 (96%)	12 (4%)	36	50
1	C	342/373 (92%)	323 (94%)	19 (6%)	21	29
1	D	346/373 (93%)	336 (97%)	10 (3%)	42	58
All	All	1377/1492 (92%)	1324 (96%)	53 (4%)	34	47

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	160	THR
1	C	255	ARG
1	D	245	ASN
1	C	192	LYS
1	C	235	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 33 ligands modelled in this entry, 10 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	C	507	-	5,5,5	0.50	0	5,5,5	0.35	0
3	H4B	D	502	-	16,18,18	0.81	0	11,26,26	2.61	6 (54%)
4	OSG	C	503	-	26,26,26	0.84	1 (3%)	35,36,36	1.41	4 (11%)
5	BTB	A	505	-	13,13,13	0.36	0	7,16,16	0.77	0
2	HEM	C	501	1	41,50,50	1.49	5 (12%)	45,82,82	1.76	10 (22%)
5	BTB	D	505	-	13,13,13	0.52	0	7,16,16	0.76	0
5	BTB	A	504	9	13,13,13	0.40	0	7,16,16	0.59	0
2	HEM	D	501	1	41,50,50	1.64	6 (14%)	45,82,82	2.08	14 (31%)
2	HEM	A	501	1	41,50,50	1.58	7 (17%)	45,82,82	1.75	12 (26%)
5	BTB	D	504	9	13,13,13	0.54	0	7,16,16	0.43	0
5	BTB	C	504	9	13,13,13	0.60	0	7,16,16	0.96	1 (14%)
4	OSG	B	503	-	26,26,26	0.88	0	35,36,36	1.33	4 (11%)
7	GOL	A	507	-	5,5,5	0.33	0	5,5,5	0.33	0
4	OSG	D	503	-	26,26,26	0.83	0	35,36,36	1.25	3 (8%)
3	H4B	A	502	-	16,18,18	1.01	0	11,26,26	2.64	6 (54%)
4	OSG	A	503	-	26,26,26	0.79	0	35,36,36	1.37	5 (14%)
3	H4B	B	502	-	16,18,18	0.90	0	11,26,26	2.72	7 (63%)
5	BTB	C	505	-	13,13,13	0.57	0	7,16,16	0.40	0
2	HEM	B	501	1	41,50,50	1.57	6 (14%)	45,82,82	2.00	13 (28%)
3	H4B	C	502	-	16,18,18	0.92	1 (6%)	11,26,26	2.77	6 (54%)
5	BTB	B	504	9	13,13,13	0.44	0	7,16,16	0.27	0
5	BTB	B	508	-	13,13,13	0.79	0	7,16,16	1.42	2 (28%)
5	BTB	B	505	-	13,13,13	0.49	0	7,16,16	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	C	507	-	-	2/4/4/4	-
3	H4B	D	502	-	-	4/8/17/17	0/2/2/2
4	OSG	C	503	-	-	0/10/19/19	0/3/3/3
5	BTB	A	505	-	-	5/21/21/21	-
2	HEM	C	501	1	-	5/12/54/54	-
5	BTB	D	505	-	-	15/21/21/21	-
5	BTB	A	504	9	-	6/21/21/21	-
2	HEM	D	501	1	-	4/12/54/54	-
2	HEM	A	501	1	-	5/12/54/54	-
5	BTB	D	504	9	-	5/21/21/21	-
5	BTB	C	504	9	-	2/21/21/21	-
4	OSG	B	503	-	-	1/10/19/19	0/3/3/3
7	GOL	A	507	-	-	4/4/4/4	-
4	OSG	D	503	-	-	3/10/19/19	0/3/3/3
3	H4B	A	502	-	-	3/8/17/17	0/2/2/2
4	OSG	A	503	-	-	1/10/19/19	0/3/3/3
3	H4B	B	502	-	-	4/8/17/17	0/2/2/2
5	BTB	C	505	-	-	5/21/21/21	-
2	HEM	B	501	1	-	4/12/54/54	-
3	H4B	C	502	-	-	3/8/17/17	0/2/2/2
5	BTB	B	504	9	-	3/21/21/21	-
5	BTB	B	508	-	-	5/21/21/21	-
5	BTB	B	505	-	-	9/21/21/21	-

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	HEM	FE-NB	4.25	2.17	1.96
2	B	501	HEM	C3C-CAC	4.20	1.56	1.47
2	D	501	HEM	C3C-CAC	4.06	1.56	1.47
2	A	501	HEM	C3C-CAC	3.97	1.56	1.47
2	D	501	HEM	C3C-C2C	-3.85	1.35	1.40

The worst 5 of 93 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	H4B	C8A-C4A-C4	5.37	119.34	114.57
3	D	502	H4B	C8A-C4A-C4	5.17	119.16	114.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	H4B	C8A-C4A-C4	5.05	119.05	114.57
3	B	502	H4B	C8A-C4A-C4	4.81	118.84	114.57
2	D	501	HEM	CBA-CAA-C2A	-4.77	104.48	112.62

There are no chirality outliers.

5 of 98 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C2A-CAA-CBA-CGA
2	A	501	HEM	C2D-C3D-CAD-CBD
2	C	501	HEM	C2A-CAA-CBA-CGA
3	A	502	H4B	C7-C6-C9-O9
3	A	502	H4B	C7-C6-C9-C10


There are no ring outliers.


21 monomers are involved in 49 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	502	H4B	1	0
4	C	503	OSG	1	0
5	A	505	BTB	4	0
2	C	501	HEM	2	0
5	D	505	BTB	3	0
5	A	504	BTB	1	0
2	D	501	HEM	4	0
2	A	501	HEM	4	0
5	D	504	BTB	2	0
5	C	504	BTB	5	0
4	B	503	OSG	2	0
7	A	507	GOL	1	0
4	D	503	OSG	2	0
3	A	502	H4B	1	0
4	A	503	OSG	2	0
5	C	505	BTB	1	0
2	B	501	HEM	4	0
3	C	502	H4B	1	0
5	B	504	BTB	1	0
5	B	508	BTB	5	0
5	B	505	BTB	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/440 (91%)	0.28	20 (4%) 28 35	31, 62, 115, 147	0
1	B	402/440 (91%)	-0.09	4 (0%) 82 86	26, 44, 81, 125	0
1	C	401/440 (91%)	-0.03	2 (0%) 91 94	30, 52, 90, 116	0
1	D	404/440 (91%)	-0.09	1 (0%) 95 96	23, 43, 74, 128	0
All	All	1611/1760 (91%)	0.02	27 (1%) 70 76	23, 50, 98, 147	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	ARG	7.2
1	A	109	LEU	4.2
1	A	106	PRO	4.0
1	A	480	TRP	3.8
1	A	204	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BTB	C	505	14/14	0.79	0.12	56,88,93,94	0
5	BTB	B	505	14/14	0.83	0.18	43,63,73,75	0
5	BTB	A	505	14/14	0.85	0.15	77,87,91,92	0
3	H4B	D	502	17/17	0.90	0.17	29,54,63,75	0
3	H4B	A	502	17/17	0.90	0.24	60,75,85,86	0
3	H4B	B	502	17/17	0.90	0.20	48,58,68,81	0
3	H4B	C	502	17/17	0.90	0.28	42,59,73,80	0
5	BTB	D	504	14/14	0.90	0.23	40,66,78,78	0
5	BTB	C	504	14/14	0.92	0.23	29,72,93,98	0
5	BTB	B	504	14/14	0.92	0.16	33,55,78,84	0
5	BTB	A	504	14/14	0.92	0.24	76,85,91,92	0
5	BTB	B	508	14/14	0.93	0.19	43,66,74,74	0
5	BTB	D	505	14/14	0.93	0.18	62,71,84,91	0
7	GOL	A	507	6/6	0.93	0.15	56,60,69,74	0
7	GOL	C	507	6/6	0.93	0.20	43,55,64,69	0
9	GD	A	509	1/1	0.93	0.12	119,119,119,119	0
4	OSG	A	503	24/24	0.94	0.14	38,59,70,70	0
4	OSG	C	503	24/24	0.94	0.14	36,50,59,61	0
8	CL	A	508	1/1	0.95	0.07	55,55,55,55	0
9	GD	C	509	1/1	0.95	0.13	99,99,99,99	0
4	OSG	B	503	24/24	0.96	0.15	19,35,63,66	0
4	OSG	D	503	24/24	0.96	0.11	17,38,57,61	0
2	HEM	A	501	43/43	0.97	0.18	43,56,81,95	0
8	CL	C	508	1/1	0.97	0.09	55,55,55,55	0
8	CL	B	506	1/1	0.98	0.11	47,47,47,47	0
2	HEM	D	501	43/43	0.98	0.14	17,34,46,65	0
8	CL	D	507	1/1	0.98	0.07	47,47,47,47	0
2	HEM	B	501	43/43	0.98	0.14	18,36,60,79	0
9	GD	B	507	1/1	0.98	0.21	53,53,53,53	0
2	HEM	C	501	43/43	0.98	0.15	27,42,70,90	0
6	ZN	A	506	1/1	0.99	0.12	49,49,49,49	0
9	GD	D	506	1/1	0.99	0.19	54,54,54,54	0
6	ZN	C	506	1/1	1.00	0.12	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around OSG B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OSG D 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.